

Oxidative functionalisation of ethane with hydrogen peroxide catalysed by chromic acid[†]

Georgiy B. Shul'pin^{a*}, Georg Süß-Fink^b and Lidia S. Shul'pina^c


^a *Semenov Institute of Chemical Physics, Russian Academy of Sciences, ul. Kosygina, dom 4, Moscow 117977, Russia*

^b *Institut de Chimie, Université de Neuchâtel, Avenue de Bellevaux 51, CH-2000 Neuchâtel, Switzerland*

^c *Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Sciences, ul. Vavilova, dom 28, Moscow 117813, Russia*

Chromic acid catalyses efficiently (turnover numbers attain 620) the oxidation of alkanes including the very inert ethane by H₂O₂ or *tert*-BuOOH in acetonitrile solution at 60 °C; alkyl hydroperoxides, ketones (aldehydes) and alcohols are the main products.

Chromium complexes are known to catalyse the oxidation of various organic derivatives (see refs 1–4) particularly C–H compounds (see refs 5–13). It should be noted that compounds containing “activated” benzylic groups^{8,10,11,13} or tertiary C–H bonds (*e.g.*, adamantane)⁹ and methylbenzenes¹² were used as substrates; molecular oxygen, peroxides, iodosobenzene were employed in these reactions as oxidising reagents. It has been shown^{14,15} that also the relatively inert cyclohexane can be oxidised by hydrogen peroxide at room temperature when a Cr(VI) oxocomplex is used as the catalyst, however the product yields were very low (the total concentration of products did not exceed 0.015 mol/dm³ after 7 days, corresponding to a turnover number, TON, of *ca* 30). On the other hand, the selective functionalisation of saturated hydrocarbons, especially of the very inert light alkanes (ethane and methane), under mild conditions remains a challenging problem for metal complex catalysis.^{5–7}



Herein we report that chromic acid catalyses the oxidation of the alkanes with hydrogen peroxide under mild conditions; the reaction is particularly efficient for the very inert ethane: CH₃CH₃ is oxidised by H₂O₂ in acetonitrile solution containing catalytic quantities of H₂CrO₄ at 60 °C to give ethyl hydroperoxide, acetaldehyde, ethanol and acetic acid (reaction 1), the total turnover number being 620 after 1 hour and the maximum yield equals 21% based on H₂O₂ (Table 1, entry 3). It follows from Table 1 that the relative content of these products depends on the initial concentration of hydrogen peroxide; at high concentrations of it, acetaldehyde becomes a predominant product. It is necessary to note that acetic acid under the conditions used is also produced via the hydrolysis of acetonitrile, however, the blank experiments demonstrated that the concentration of CH₃COOH formed from CH₃CN was only 0.002 mol/dm³. Propane under analogous conditions (but at lower pressure, 5 bar) gave (concentrations are indicated in parentheses, mol/dm³) CH₃CH₂CHO (0.009), CH₃COCH₃ (0.026), CH₃CH(OH)CH₃ (0.031), CH₃CH₂CH₂OH (0.02) and CH₃CH₂COOH (0.009) (TON 190). Butane (atmospheric pressure, 1 bar) gave CH₃(CH₂)₂CHO (0.0003), CH₃COCH₂CH₃

(0.0052), CH₃CH(OH)CH₂CH₃ (0.012), CH₃(CH₂)₂CH₂OH (0.0011) and CH₃(CH₂)₂COOH (0.00037) (TON 38).

The oxidations of ethane and other gaseous alkanes were carried out at 60 °C in a stainless steel autoclave (the volume of the reaction solution being 10 ml, the total autoclave volume being 100 ml). The oxidations of higher hydrocarbons were carried out in air in thermostated (60 °C) Pyrex cylindrical vessels with vigorous stirring. The total volume of the reaction solution was 10 ml. In all experiments, a 35% aqueous solution of hydrogen peroxide (“Fluka”) or 70% aqueous solution of *tert*-butyl hydroperoxide (“Fluka”) was used. The catalyst, H₂CrO₄, usually employed in concentration 5.0 × 10⁻⁴ mol/dm³, was introduced into the reaction mixture in the form of a stock solution (0.1 ml), which has been prepared by dissolving CrO₃ (0.025 g) in H₂O (5 ml). The samples of the reaction solutions were analysed by GC (DANI-86.10; fused silica capillary column) twice: before and after addition of an excess of solid triphenylphosphine. Triphenylphosphine reduces hydrogen peroxide to water and the alkyl hydroperoxide to the corresponding alcohol which allowed the determination of the *real* concentrations of the alkyl hydroperoxide, formed from the alkane, as well as those of the alcohol and the ketone (or aldehyde). In some cases, it is possible to find peaks corresponding to alkyl hydroperoxides. These peaks disappear completely after the treatment of the solution with PPh₃, while peaks of the corresponding alcohols grow (see ref. 19). We have developed and used this method^{5,6,16–21} for the analysis of reaction mixtures obtained from various alkane oxidations.

In order to get a mechanistic understanding of this reaction, we also studied the oxidation of higher alkanes with hydrogen peroxide in the presence of chromic acid. The main product of cyclooctane oxidation was the alkyl hydroperoxide which gradually decomposed to give predominantly cyclooctanone. The reaction was accelerated but insignificantly (*ca* 15%) if pyridine was added (see ref. 22). After 8 hours the turnover number attained 280. During the oxidation with *t*-BuOOH, the concentration of cyclooctyl hydroperoxide was low and the reaction mixture contained cyclooctanone produced with high selectivity. In this case the rate of the reaction and the product yields were somewhat lower if pyridine was added. The oxidation with H₂O₂ in CH₃COOH as a solvent is less efficient (0.011 and 0.024 mol/dm³ of cyclooctanone and cyclooctanol, respectively, after 7 hours at 60 °C and after treatment with PPh₃).

The oxidation of *n*-octane in CH₃CN under the same conditions with H₂O₂ in the absence of pyridine gave after 3 h (after reduction with PPh₃; the concentrations are indicated in parentheses, *c* × 10² mol/dm³) octanal (0.17), octanone-1 (0.39),

* To receive any correspondence. E-mail: Shulpin@chph.ras.ru; WEB: <http://shulpin.newmail.ru>

[†] This is a Short Paper, there is therefore no corresponding material in *J. Chem. Research (M)*.

Table 1 Ethane oxidation with H₂O₂ in CH₃CN catalysed by H₂CrO₄^a

Entry	[H ₂ O ₂] ₀ , mol/dm ³	Products, mol/dm ³				
		CH ₃ CH ₂ OOH	CH ₃ CHO	CH ₃ CH ₂ OH	CH ₃ COOH	TON ^b
1	0.6	0.09	0.003	0.01	0	206
2	1.2	0.08	0.09	0.06	0.02	500
3	2.4	0.07	0.16	0.06	0.02	620

^aConditions: ethane, 30 bar; H₂CrO₄, 5 × 10⁻⁴ mol/dm³; 1 hour at 60°C. The concentrations of reaction products were measured before and after reduction with PPh₃ to determine *real* concentrations of ethyl hydroperoxide, ethyl alcohol and acetaldehyde.
^bTurnover number, *i.e.*, total number of moles of all products per one mole of the catalyst.

Table 2 Selectivities of alkane oxidations by "ROOH – H₂CrO₄" and some other systems in MeCN^a

Entry	System	Alkane oxidised (the selectivity parameter ^b)		
		<i>cis</i> -Decalin (<i>t/c</i>)	<i>trans</i> -Decalin (<i>t/c</i>)	Adamantane (3°: 2°)
1	H ₂ O ₂ – H ₂ CrO ₄ (60 °C)	3.9	4.3	1.4
2	<i>t</i> -BuOOH – H ₂ CrO ₄ (50 °C)	4.3	6.1	4.0
3	H ₂ O ₂ – <i>hν</i> (20 °C)	1.3	2.7	0.9
4	<i>t</i> -BuOOH – <i>hν</i> (20 °C)	5.1	4.4	1.1
5	H ₂ O ₂ – VO ₃ ⁻ – PCA ^c	2.1	2.4	0.7
6	H ₂ O ₂ – Mn(IV) ^d – CH ₃ COOH	0.12	33.4	2.0

^aThe concentrations of reaction products were measured after reduction with PPh₃. ^bParameter *t/c* = [*trans*-decal-9-ol]/[*cis*-decal-9-ol], *i.e.*, the ratio of concentrations of *trans*-decal-9-ol and *cis*-decal-9-ol formed in the oxidation of isomeric decalins. Parameter 3°: 2° is normalised (*i.e.*, calculated taking into account the number of hydrogen atoms at each position) relative reactivities of hydrogen atoms at tertiary and secondary carbons, respectively. ^cPCA, pyrazine-2-carboxylic acid. For this system, see refs 16, 19.
^dComplex LMn^{IV}(O)₃Mn^{IV}L, where L = 1,4,7-trimethyl-1,4,7-triazacyclononane. For this system, see refs 20, 21.

octanone-3 (0.44), octanone-2 (0.36), octanol-4 (1.65), octanol-3 (1.8), octanol-2 (2.0), octanol-1 (0.5). Taking into account only concentrations of alcohols, we can calculate the non-normalised ratio C(1) : C(2) : C(3) : C(4), *i.e.*, relative reactivities of hydrogens at 1, 2, 3 and 4 carbon atoms of the alkane chain, respectively: 1.0 : 4.0 : 3.6 : 3.3. This selectivity is higher than that obtained by us for the oxidation with the system H₂O₂ – *hν* (1.0 : 2.8 : 2.9 : 2.7)

Selectivity parameters for the oxidation of decalin (*cis*- and *trans*-isomers) and adamantane are given in Table 2. It can be seen that selectivities for the two Cr-containing systems (entries 1 and 2) are different from those obtained for HO-radical-generating systems (entries 3 and 5) and from the system capable of generating HO and *t*-BuO radicals (entry 4). It should be noted, that the Cr-containing systems oxidise non-stereoselectively unlike the Mn(IV)-based system recently described (entry 6). Thus one can conclude that the Cr-catalysed reactions does not seem to proceed via HO radicals but involve possibly peroxo complexes^{23–25} as oxidising species.

We thank the Russian Basic Research Foundation and the Swiss National Science Foundation for support. The authors are indebted to Dr Yuriy N. Kozlov for useful discussions and to Mrs Sandrine Stanislas, Mr Matthieu Faure and Mr Christophe Thomas for help in this work.

Received 11 July 2000; accepted 6 October 2000
Paper 00/445

References

- J. Muzart, *Chem. Rev.*, 1992, **92**, 113–140.
- W. Adam, F.G. Gelalcha, C.R. Saha-Möller and V.R. Stegmann, *J. Org. Chem.*, 2000, **65**, 1915–1918.
- P.A. Shapley, N. Zhang, J.L. Allen, D.H. Pool and H.-C. Liang, *J. Am. Chem. Soc.*, 2000, **122**, 1079–1091.
- A. Bakac and W.-D. Wang, *Inorg. Chim. Acta*, 2000, **297**, 27–35.
- A.E. Shilov and G.B. Shul'pin, *Chem. Rev.* 1997, **97**, 2879–2932.
- A.E. Shilov and G.B. Shul'pin, *Activation and Catalytic Reactions of Saturated Hydrocarbons in the Presence of Metal Complexes*, Kluwer Academic Publishers, Dordrecht/Boston/London, 2000.
- G.B. Shul'pin, *Organic Reactions Catalysed by Metal Complexes*, Nauka, Moscow, 1988.
- G.B. Shul'pin and A. N. Druzhinina, *Bull. Acad. Sci. USSR, Div. Chem. Sci.*, 1989, **38**, 1079–1081.
- J. Muzart, and A. N' Ait Ajjou, *J. Mol. Catal.*, 1993, **84**, L15–L19.
- J. Muzart, and A. N' Ait Ajjou, *J. Mol. Catal.*, 1994, **92**, 141–147.
- J. Muzart, and A. N' Ait Ajjou, *J. Mol. Catal.*, 1994, **92**, 277–283.
- I.C. Chisem, J. Rafelt, M.T. Shieh, J. Chisem, J.H. Clark, R. Jachuck, D. Marcquarrie, C. Ramshaw and K. Scott, *Chem. Commun.*, 1998, 1949–1950.
- N. Delaval, S. Bouquillon, F. Hémin, and J. Muzart, *J. Chem. Res. (S)*, 1999, 286–287.
- A.N. Druzhinina, G.V. Nizova and G.B. Shul'pin, *Bull. Acad. Sci. USSR, Div. Chem. Sci.*, 1990, **39**, 194–195.
- J. Muzart, A. N' Ait Ajjou, G. V. Nizova and G. B. Shul'pin, *Izvest. Akademii Nauk SSSR, Ser. Khim.*, 1991, 1454–1457.
- G.B. Shul'pin, D. Attanasio and L. Suber, *J. Catal.*, 1993, **142**, 147–152.
- G.B. Shul'pin, M.M. Bochkova and G.V. Nizova, *J. Chem. Soc., Perkin Trans. 2*, 1995, 1465–1469.
- G.B. Shul'pin, G.V. Nizova and Yu. N. Kozlov, *New. J. Chem.*, 1996, **20**, 1243–1256.
- G.B. Shul'pin, M.C. Guerreiro and U. Schuchardt, *Tetrahedron*, 1996, **52**, 13051–13062.
- G.B. Shul'pin and J. R. Lindsay Smith, *Russ. Chem. Bull.*, 1998, **47**, 2379–2386.
- G.B. Shul'pin, G. Süß-Fink and J.R. Lindsay Smith, *Tetrahedron*, 1999, **55**, 5345–5358.
- G. Rothenberg, H. Wiener and Y. Sasson, *J. Mol. Catal., A: Chem.*, 1998, **136**, 253–262.
- E. Daire, H. Mimoun and L. Saussine, *Nouv. J. Chim.*, 1984, **8**, 271–274.
- A. Bakac, T.-J. Won and J. H. Espenson, *Inorg. Chem.*, 1996, **35**, 2171–2175.
- U. El-Ayaan, R. Reisacher and W. Linert, *J. Chem. Res. (S)*, 1999, 521.